

HOW TO MANAGE MY ASTHMA AFTER DISCHARGE FROM HOSPITAL

Most children with asthma and many children with pre-school wheeze will be given a course of steroids (prednisolone) when they become unwell.

- If you have been started on a course of prednisolone, it is important that you complete the course that you have been given.
- Give the reliever inhaler via spacer as needed, up to 10 puffs every 4 hours
- Keep the regular treatments going
- Avoid over-excitement
- Please check on your child overnight
- If your child needs more reliever than 10 puffs 4 hourly see your GP today or visit A/E
- You should make an appointment to see your GP within 48 hours of discharge

GP ASTHMA CLINIC

Ask for a review if

- Your child is getting lots of mild attacks
- Your child starts using their reliever inhaler more than 3 times a week.
- Use the "Asthma Control Test" to see how you are doing. Find it at www.asthma.org.uk

BEFORE EXERCISE

Taking my reliever before I start may help prevent me wheezing and coughing.

TRIGGERS THAT MAKE ME WHEEZE

Please list below:

THESE ARE MY CONTACTS

Doctor (GP):

Nurse:

Hospital consultant:



Want to give
up smoking?

For free help and advice see your
G.P or call for free

0800 085 2219

www.helpmequit.wales



ASTHMA & PRE-SCHOOL WHEEZE

Management Plan

Patient Addressograph

Last updated:

Learn more
more here



Y Grŵp Gweithredu
ar Iechyd Anadiol
Respiratory Health
Implementation Group

Iechyd Plant

Child Health



Generated from
Asthmahub for Parents

MONITORING YOUR ASTHMA

IF YOUR CHILD IS GETTING WORSE

- Wheeze and coughing is getting worse
- Your child is feeling out of breath

WHAT TO DO NEXT

 Increase reliever inhaler to 10 puffs every 4 hours via spacer

 Continue your preventer treatment

 See your GP today or attend A&E

 Give Home Rescue Steroid if directed to do so

IF YOUR CHILD IS HAVING A SEVERE ATTACK?!

- Distressed by cough and wheeze or breathing hard?
- Won't play because of breathlessness
- Is too breathless to speak
- Reliever therapy doesn't last long

 This is life threatening: Act early: Call 999 or go to A&E immediately

WHILST WAITING TO BE SEEN!

- Give 10 puffs of the reliever inhaler via spacer
- Continue with 10 puffs reliever via spacer as often as needed
- If you have Home Rescue Steroid give a dose as soon as possible

IF YOUR CHILD...

- Needs reliever inhaler more than usual
- Is coughing or wheezing more
- Is coughing and waking at night

WHAT TO DO NEXT

 Increase reliever inhaler to 6-10 puffs every 4 hours via spacer

 If not improving, see your GP today

 Continue your preventer treatment

 If improving see your GP in the next few days

WHEN YOUR CHILD IS WELL

 Give the preventer treatment as prescribed

SYMPTOM CHECKER

AS YOUR CHILD IMPROVES

Give up to 10 puffs of reliever via spacer as needed

Seek advice if 10 puffs doesn't last 4 hours

Check on your child overnight

REGULAR TREATMENT

Reliever therapy

Preventer therapy

Home rescue steroid

I ALWAYS USE

- Spacer and mouthpiece
- Spacer and mask

Mouthpiece spacers are far more effective than mask spacers. Children age >3years should be able to use a mouthpiece spacer

WHEN I'M UNWELL, MY WHEEZING

- Responds to steroids
- Does not respond to steroids
- We are not sure